

ATTORNEY'S DOCKET NO: 24406

U.S. DEPARTMENT OF COMMERCE, PATENT AND TRADEMARK OFFICE		DATE: <u>16</u> October 2000 (<u>16</u> 10.2000)
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		U.S. PAT. NO. (if known): Not Yet Assigned 09/673473
INTERNATIONAL APPLICATION NO.: PCT/CA99/00311	INTERNATIONAL FILING DATE: 19 April 1999 (19.04.99)	PRIORITY DATE CLAIMED: 17 April 1998 (17.07.98)
TITLE OF INVENTION: CUBANE DERIVATIVES AS METABOTROPIC GLUTAMATE RECEPTOR AGONISTS OR ANTAGONISTS AND PROCESS FOR THEIR PREPARATION		
APPLICANT(S) FOR DO/EO/US: CURRY, Kenneth; PAJOUHESH, Hassan		
Applicant hereby submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:		
<ol style="list-style-type: none"> <u>X</u> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. <u> </u> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. <u>X</u> This express request to begin national examination procedures (35 USC 371(f)) at any time rather than delay examination until the expiration of the time limit set in 35 USC 371(b) and PCT Articles 22 and 39(1). <u>X</u> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. <u>X</u> A copy of the International Application as filed (35 U.S.C. 371(c)(2)): <ol style="list-style-type: none"> <u>X</u> is transmitted herewith (required only if not transmitted by the International Bureau). <u> </u> has been transmitted by the International Bureau. <u> </u> is not required, as the application was filed in the United States Receiving Office (RO/US) <u> </u> A translation of the International Application into English (35 U.S.C. 371(c)(2)). <u>X</u> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> <u> </u> are transmitted herewith (required only if not transmitted by the International Bureau). <u> </u> have been transmitted by the International Bureau. <u> </u> have not been made; however, the time limit for making such amendments has NOT expired. <u>X</u> have not been made and will not be made. <u> </u> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). <u> </u> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). <u> </u> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). 		
ITEMS 11. TO 16. BELOW CONCERN OTHER DOCUMENT(S) OR INFORMATION INCLUDED:		
<ol style="list-style-type: none"> <u> </u> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. <u> </u> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. <u>X</u> A FIRST preliminary amendment. <u> </u> A SECOND or SUBSEQUENT preliminary amendment <u> </u> A substitute specification. <u> </u> A change of power of attorney and/or address letter. <u>X</u> TRANSMITTAL FORM; FEE CALCULATION; INTERNATIONAL PUBLICATION WO99/54280; INTERNATIONAL PUBLICATION DATE 28 OCTOBER 1999; APPLICATION CONSISTING OF 42 PAGES INCLUDING; 36 PAGES TEXTUAL SPECIFICATION, 4 PAGES OF 10 CLAIMS; 1 COVER SHEET CONTAINING THE ABSTRACT; 1 SHEET DRAWINGS; PRELIMINARY AMENDMENT; UNEXECUTED INVENTOR'S DECLARATION; PCT/IPEA/408 WRITTEN OPINION; RESPONSE TO WRITTEN OPINION DATED JULY 19, 2000; PCT/ISA/210 INTERNATIONAL SEARCH REPORT; PCT/RO/101 REQUEST/ PCT/IPEA/401 DEMAND; PCT/IPEA/402 NOTIFICATION OF RECEIPT OF DEMAND BY COMPETENT INTERNATIONAL PRELIMINARY EXAMINING AUTHORITY. 		

ATTORNEY'S DOCKET NO: 24406

[illegible]

ATTORNEY'S DOCKET NO: 24406

U.S. APPLICATION NO. Not yet assigned 09/673473	INTERNATIONAL APPLICATION NO. PCT/CA99/00311	DATE: 16 October 2000 (16 .10.2000)
--	--	--

a. ☒ One check in the amount of \$878.00 to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account No. 14-0112 in the amount of \$_____ to cover the above fees. (A duplicate copy of this sheet is enclosed.)


c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 14-0112.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed to request that the application be restored to pending status.

Send All Correspondence To:

Gary M. Nath
NATH & ASSOCIATES PLLC
1030 15th Street, N.W.
Sixth Floor
Washington, D.C. 20005

(202) 775-8383 (phone)
(202) 775-8396 (fax)



GARY M. NATH
Registration Number 26,965
Customer No. 20529

Rev. 02/98

Rec'd PCT/PTO 9 NOV 2000

526 Rec'd PCT/PTO 2 NOV 2000

BOX PCT-MISSING REQUIREMENTS
Attorney Docket No. 24406

12/6/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

CURRY, Kenneth; PAJOUHESH, Hassan


International Application No. PCT/CA99/00311

Serial No. 09/673,473

International Filing Date: 19 April 1999 (19.04.99)

Filed: October 16, 2000

**CUBANE DERIVATIVES AS METABOTROPIC GLUTAMATE RECEPTOR
AGONISTS OR ANTAGONISTS AND PROCESS FOR THEIR PREPARATION**



TRANSMITTAL LETTER

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Submitted herewith for filing in the U.S. Patent and
Trademark Office is the following:

- (1) Transmittal Letter
- (2) Copy of Notification of Missing Requirements Under 35 U.S.C.
371
- (3) Response to Notification of Missing Requirements
- (4) Executed Declaration and Power of Attorney
- (5) Assertion that Applicant is entitled to claim small entity
status
- (6) Request for Refund
- (7) Copies of Check No. 13568, stamped filing receipt and post
card showing payment of filing fee of \$878.00
- (8) Check No. 13859 \$ 65.00 surcharge for late filing of
Declaration

The Commissioner is hereby authorized to charge any
deficiency or credit any excess to Deposit Account No. 14-0112.

Respectfully submitted,

11/30/2000 ERIMANDO 00000144 09673473

01 FC:254

65.00 DP

NATH & ASSOCIATES PLLC

By: _____

Gary M. Nath
Registration No. 26,965
Customer No. 20529

Date: November 29, 2000

NATH & ASSOCIATES PLLC

1030 15th Street N.W., 6th Floor

Washington, D.C. 20005

(202) 775-8383

GMN/dd:NEMP.TRANS

PATENT
Attorney Docket No. 24406

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Kenneth CURRY et al.

International Application No. PCT/CA99/00311

Serial No: (Not Yet Assigned)

International Filing Date: 19 April 1999 (10.04.99)

Filed: October 16, 2000

For: CUBANE DERIVATIVES AS METABOTROPIC GLUTAMATE RECEPTOR
AGONISTS OR ANTAGONISTS AND PROCESS FOR THEIR PREPARATION

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, D.C. 20231

Sir:

Before action in this application and before calculation of the filing fee, please amend the above identified application as follows:

IN THE TITLE

In the title, please replace:

"Cubane Derivatives as Metabotropic Glutamate Receptor Antagonists and Process for their Preparation"

with:

--Cubane Derivatives as Metabotropic Glutamate Receptor Agonists or Antagonists and Process for their Preparation--.


IN THE SPECIFICATION

Page 8, line 7, after "R2 is", please replace "COOH" with --R2 is NH₂--.

Page 8, line 8, after "xanthyl or thioxanthyl" and before "and", please insert --or -CH₂-xanthyl or -CH₂-thioxanthyl--.

Page 8, line 9, after "**R4** is", please replace "NH₂" with --COOH--.


Page 11, lines 5-9, please replace:
 "(a) hydrolyzing a compound of formula:



in which **R1** is defined as above, **R5** represents a hydrogen atom or an acyl group and **R4** has the meaning defined above. Preferred values for **R5** are hydrogen and (2-6C) alkanoyl groups, such as acetyl."

with:

--(a) hydrolyzing a compound of formula (IIa):




wherein: **R'1** is an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono. Sulfinio, borono, tetrazol, isoxazol, -CH₂-carboxyl, -CH₂-phosphono, -CH₂-phosphino, -CH₂-sulfono, -CH₂-sulfinio, -CH₂-borono, -CH₂-tetrazol, -CH₂-isoxazol and higher analogues thereof, or a protected form thereof, **R3** can be H, aliphatic, aromatic or heterocyclic

and **R5** represents a hydrogen atom or an acyl group. Preferred values for **R5** are hydrogen and (2-6C) alkanoyl groups, such as acetyl; or--.

Page 11, lines 8-9, after "Preferred values for **R5** are hydrogen and (2-6C) alkanoyl groups, such as acetyl; or", please insert:


--(b) deprotecting and hydrolyzing a compound of formula (IIb)



wherein: **R'1** and **R3** are as defined above; or--.

Page 11, lines 11-15, please remove:

"(b) hydrolyzing a compound of formula:




in which **R6** and **R7** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is

unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof; or:"

and insert on new page 11a:


--(c) hydrolyzing a compound of formula:



wherein: **R'1** and **R3** has the meaning defined above, **R6** and **R7** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof; or:-- .

Page 12, lines 1-4, please replace:


"(c) deprotecting a compound of formula:



in which **R8** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R9** represents a hydrogen atom or a nitrogen protecting group;"

with:

--(d) deprotecting a compound of formula:




wherein: **R'1** and **R3** has the meaning defined above, **R8** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R9** represents a hydrogen atom or a nitrogen protecting group;--.

Page 13, line 17, after "heating the compound of" and before "in the presence of either a base,", please replace "Formula V" with --Formula IV--.

Page 13, line 22, after "reacting the compound of" and before "with hydrogen in the presence", please replace "Formula V" with --Formula IV--.


Page 13, lines 29-30, please replace:

"The compounds of Formula II may be prepared by reacting a compound of formula:




with:

--The compounds of Formula II may be prepared by reacting a compound of formula (V):



Page 14, lines 20-22, please replace:


"The compounds of Formula V can be prepared by reacting a compound of formula:



with a chlorinating agent such as thionyl chloride or phosphorous (V) chloride, followed by reaction with $R4X$ wherein $R4$ has the meaning defined above and X is halogen or OH."

with:


--The compounds of Formula V can be prepared by reacting a compound of formula:



with a chlorinating agent such as thionyl chloride or phosphorous (V) chloride, followed by reaction with organo copper or organo metal or Grignard reagent derived from $R3X$ or by reaction with ethyl hydrogen malonate in the presence of organolithium, wherein $R3$ has the meaning defined above and X is halogen.--.


Page 15, lines 1-10, please replace:

"The compounds of Formula V can also be prepared by oxidizing a compound of formula



under Swern conditions.

The compounds of Formula VI can be prepared from compounds of formula:




by reduction.

If **R¹** is CO₂Me, this compound can be bought commercially. If **R¹** is another substituent, the compound of Formula VIII can be made using standard procedures."


with:

--The compounds of Formula V can also be prepared by oxidizing a compound of formula:



under Swern conditions.

The compounds of Formula VI can be prepared from compounds of formula:



by reduction.


When **R'1** is CO₂Me, this compound can be bought commercially. When **R'1** is another substituent, the compound of Formula VIII can be made using standard procedures.--.

Page 31, line 26, please replace "*Preparation 5: 4-carbonyl-1-cubanylglycine*" with --*Preparation 5: 4-carboxy-1-cubanylglycine*--.


IN THE CLAIMS

5. (Amended) A process for the preparation of a compound of Formula I, or a pharmaceutically acceptable metabolically-labile ester or amide thereof, or a pharmaceutically acceptable salt thereof, which comprises:

(a) hydrolyzing a compound of formula: [




in which **R1** is defined as above,]




wherein: **R'1** is an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol, -CH₂-carboxyl, -CH₂-phosphono, -CH₂-phosphino, -CH₂-sulfono, -CH₂-sulfino, -CH₂-borono, -CH₂-tetrazol, -CH₂-isoxazol and higher analogues thereof, or a protected form thereof, **R3** can be H, aliphatic, aromatic or heterocyclic and **R5** represents a hydrogen atom or an acyl group [and **R4** has the meaning defined above.], and wherein preferred [Preferred] values for **R5** are hydrogen and (2-6C) alkanoyl groups, such as acetyl;
or

(b) deprotecting and hydrolyzing a compound of formula (IIb):





wherein: **R'1** and **R3** are as defined above; or

(c) [(b)] hydrolyzing a compound of formula:



wherein: **R6** and **R7** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof, **R'1** and **R3** are as defined above; or

(d) [(c)] deprotecting a compound of formula:



wherein: [in which] **R8** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R9** represents a hydrogen atom or a nitrogen protecting group, **R'1** and **R3** are as defined above;



whereafter, if necessary and/or desired:

- (i) resolving the compound of Formula I;
- (ii) converting the compound of Formula I into a non-toxic metabolically-labile ester or amide thereof; and/or[;]
- (iii) converting the compound of Formula I or a non-toxic metabolically-labile ester or amide thereof into a pharmaceutically acceptable salt thereof.

7. (Amended) A use of the compound according to claim 1 to modulate [A method of modulating] one or more metabotropic


glutamate receptor functions in a warm blooded mammal [requiring such treatment], wherein said use [which] comprises administering an effective amount of a compound of formula (I) as claimed in claim 1.

8. (Amended) A compound of formula:





wherein: [in which **R1**, **R4**] **R'1**, **R3** and **R5** have the meanings as defined in claim 5 [above].

9. (Amended) A compound of formula:



wherein: R'1, R3, [wherein] R6 and R7 have meanings as defined in claim 5 [above].

10. (Amended) A compound of formula:



wherein: R'1, R3, [in which] **R8** and **R9** have meanings as defined in claim 5 [above].

Please add the following newly submitted claims.

--11. A compound according to claim 1, wherein **R1** is -COOH, **R2** is -NH₂, **R3** is H and **R4** is COOH.

12. A compound according to claim 1, wherein **R1** is -COOH, **R2** is -NH₂, **R3** is CH₃ and **R4** is COOH.


13. A compound according to claim 1, wherein **R1** is -COOH, **R2** is -NH₂, **R3** is -CH₂-thioxanthy and **R4** is COOH.

14. A use of the compound according to claim 1 for the treatment of a neurological disease or disorder selected from the group comprising: cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia, stroke, cardiac arrest,


spinal cord trauma, head trauma, perinatal hypoxia, and hypoglycemic neuronal damage, Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, AIDS-induced dementia, ocular damage, retinopathy, cognitive disorders, idiopathic and drug-induced Parkinson's disease, muscular spasms, convulsions, migraine headaches, urinary incontinence, psychosis, drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol), smoking cessation, anxiety and related disorders (e.g. panic attack), emesis, brain edema, chronic pain, sleep disorders, Tourette's syndrome, attention deficit disorder, and tardive dyskinesia, wherein said use comprises administering an effective amount of a compound of formula (I).

15. A use of the compound according to claim 1 for the treatment of a psychiatric disease or disorder selected from the group comprising: schizophrenia, anxiety and related disorders (e.g. panic attack), depression, bipolar disorders, psychosis, and obsessive compulsive disorders, wherein said use comprises administering an effective amount of a compound of formula (I).

16. The use according to claim 7 wherein said compound is selected from the group of compounds comprising:




17. A use of the compound:



for the treatment of cerebral ischemia, stroke and cardiac arrest, wherein said use comprises administering an effective amount of the said compound.

18. A compound of formula:



wherein: **R'1** and **R3** have the meaning as defined in claim 5.

19. A compound according to claim 18, wherein: **R'1** is -COOMe, **R3** is H.

20. A compound according to claim 9, wherein: **R'1** is -COOH, **R3** is CH₃, **R6** = **R7** is H.

21. A compound according to claim 9, wherein: **R'1** is -COOH, **R3** is -CH₂-thioxanthyl, **R6** = **R7** is H.--

REMARKS

Upon entry of the above amendments, claims 1-21 are pending in the application. The claims and specification have been amended in accordance with the Article 34 Amendments submitted in the corresponding International Application No. PCT/CA99/00311. In particular, the specification was amended to correct typographical and clerical errors. The amendments do not introduce new matter within the meaning of 35 U.S.C. § 132. Accordingly, entry of the amendments is respectfully requested.

CONCLUSION


In view of the foregoing, Applicants respectfully request the Examiner to allow all claims pending in this application. If

the Examiner has any questions or wishes to discuss this matter,
the Examiner is welcomed to telephone the undersigned attorney.

Respectfully submitted,

NATH & ASSOCIATES PLLC

Date: October 16, 2001



Gary M. Nath
Reg. No. 26,965
Customer No. 20529

NATH & ASSOCIATES PLLC
1030 Fifteenth Street, N.W.
Sixth Floor
Washington, D.C. 20005-1503
Tel: (202) 775-8383
Fax: (202) 775-8396

GMN:JLM:JBG:\PRELIM.wpd

CUBANE DERIVATIVES AS METABOTROPIC GLUTAMATE RECEPTOR ANTAGONISTS AND PROCESS FOR THEIR PREPARATION

FIELD OF THE INVENTION

This invention pertains to therapeutically active cubane derivatives, a method for preparing the same, pharmaceutical compositions comprising the compounds and a method of treating diseases of the Central Nervous System (CNS) therewith.

BACKGROUND OF THE INVENTION

The acidic amino acid L-Glutamate is recognized as the major excitatory neurotransmitter in the CNS. The receptors that respond to L-Glutamate are called excitatory amino acid receptors. The excitatory amino acid receptors are thus of great physiological importance, playing a role in a variety of physiological processes, such as long-term potentiation (learning and memory), the development of synaptic plasticity, motor control, respiratory and cardiovascular regulation, and sensory perception.

Excitatory amino acid receptors are classified into two general types and both are activated by L-Glutamic acid and its analogs. Receptors activated by L-Glutamic acid that are directly coupled to the opening of cation channels in the cell membrane of the neurons are termed "ionotropic." This type of receptor has been subdivided into at least three subtypes, which are defined by the depolarizing actions of the selective agonists N-Methyl-D-aspartate (NMDA), α -Amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA), and Kainic acid (KA).

The second general type of receptor is the G-protein or second messenger-linked "metabotropic" excitatory amino acid receptor. This second type is coupled to multiple second messenger systems that lead to enhanced phosphoinositide hydrolysis, activation of phospholipase D, increases or decreases in cAMP formation, and changes in ion channel function (Schoepp and Conn, *Trends in Pharmacological Science*, 14:13, 1993). Both types of receptors appear not only to mediate normal synaptic transmission along excitatory pathways but also to participate in the modification of synaptic connections during development and throughout life.

So far eight different clones of the G-protein-coupled metabotropic glutamate receptors (mGluRs) have been identified (Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426). These receptors function to modulate the presynaptic release of L-Glutamate, and the postsynaptic

sensitivity of the neuronal cell to L-Glutamate excitation. Based on pharmacology, sequence homology and the signal transduction pathway that they activate, the mGluRs have been subclassified into three groups. The mGluR1 and mGluR5 receptors form group I. They are coupled to hydrolysis of phosphatidylinositol (PI) and are selectively activated by (*RS*)-3,5-dihydroxyphenylglycine (Brabet et al., *Neuropharmacology*, 34, 895-903, 1995). Group II comprises mGluR₂ and mGluR₃ receptors. They are negatively coupled to adenylate cyclase and are selectively activated by (*2S,1'R,2'R,3'R*)-2-(2,3-dicarboxycyclopropyl)glycine (DCG-IV; Hayashi et al., *Nature*, 366, 687-690, 1993). Finally, the mGluR₄, mGluR₆, mGluR₇ and mGluR₈ receptors belong to group III. They are also negatively coupled to adenylate cyclase and are selectively activated by (L)-2-amino-4-phosphonobutyric acid (L-AP4; Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426).

Agonists and antagonists of these receptors are believed useful for the treatment of acute and chronic neurodegenerative conditions, and as antipsychotic, anticonvulsant, analgesic, anxiolytic, antidepressant, and anti-emetic agents. Antagonists and agonists of neural receptors are classified as selective for a particular receptor or receptor subtype, or as non-selective. Antagonists may also be classified as competitive or non-competitive. While competitive and non-competitive antagonists act on the receptors in a different manner to produce similar results, selectivity is based upon the observations that some antagonists exhibit high levels of activity at a single receptor type, and little or no activity at other receptors. In the case of receptor-specific diseases and conditions, the selective agonists and antagonists are of the most value.

Compounds such as L-Glutamic acid, Quisqualic acid and Ibotenic acid are known to act as non-selective agonists on the mGluRs, while selective ionotropic glutamate receptor agonists such as NMDA, AMPA and Kainic acid have little effect on these receptors. Recently a few compounds without activity at the ionotropic glutamate receptors but with activity at the metabotropic receptors have been identified. These include *trans*-ACPD (*trans* (1*S*,3*R*-1-aminocyclopentane-1,3-dicarboxylic acid), the partial agonist L-AP3 (L-2-amino-3-phosphonopropionic acid; Palmer, E., Monaghan, D. T. and Cotman, C. W. *Eur. J. Pharmacol.* 166, 585-587, 1989; Desai, M. A. and Conn, P. J. *Neuroscience Lett.* 109, 157-162, 1990; Schoepp, D. D. et al., *J. Neurochemistry*. 56, 1789-1796, 1991; Schoepp D. D. and Johnson B. G. *J. Neurochemistry* 53, 1865-1869, 1989), L-AP4 (L-2-amino-4-phosphonobutyric acid) which is an agonist at the mGluR₄ receptor (Thomsen C. et al., *Eur. J. Pharmacol.* 227, 361-362, 1992) and some of the isomers of CCG (2-(carboxycyclopropyl)glycines) especially L-CCG-I and L-CCG-II (Hayashi, Y. et al., *Br. J. Pharmacol.* 107, 539-543, 1992).

Very few selective antagonists at the mGluRs have been reported. However some phenylglycine derivatives, *S*-4CPG (*S*-4-carboxyphenylglycine), *S*-4C3HPG (*S*-4-carboxy -3-hydroxyphenylglycine) and *S*-MCPG (*S*- α -methyl-4-carboxyphenylglycine) have been reported to antagonize *trans*-ACPD-stimulated phosphoinositide hydrolysis and thus possibly act as antagonists at mGluR₁ and mGluR₅ subtypes (Thomsen, C. and Suzdak, P, *Eur. J. Pharmacol.* 245, 299, 1993).

Research directed towards mGluRs is beginning to show that mGluRs may be implicated in a number of normal as well as pathological mechanisms in the brain and spinal cord. For example, activation of these receptors on neurons can: influence levels of alertness, attention and cognition; protect nerve cells from excitotoxic damage resulting from ischemia, hypoglycemia and anoxia; modulate the level of neuronal excitation; influence central mechanisms involved in controlling movement; reduce sensitivity to pain; reduce levels of anxiety.

The use of compounds active at the mGluRs for the treatment of epilepsy is corroborated by investigations of the influence of *trans*-ACPD on the formation of convulsions (Sacaan and Schoepp, *Neuroscience Lett.* 139, 77, 1992) and that phosphoinositide hydrolysis mediated via mGluR is increased after kindling experiments in rats (Akiyama et al. *Brain Res.* 569, 71, 1992).

Trans-ACPD has been shown to increase release of dopamine in the rat brain, which indicates that compounds acting on the mGluRs might be usable for the treatment of Parkinson's disease and Huntington's Chorea (Sacaan et al., *J. Neurochemistry* 59, 245, 1992).

Trans-ACPD has also been shown to be a neuroprotective agent in a medial cerebral artery occlusion (MCAO) model in mice (Chiamulera et al. *Eur. J. Pharmacol.* 215, 353, 1992), and it has been shown to inhibit NMDA-induced neurotoxicity in nerve cell cultures (Koh et al., *Proc. Natl. Acad. Sci. USA* 88, 9431, 1991). The mGluR-active compounds are also implicated in the treatment of pain. This is proved by the fact that antagonists at the metabotropic glutamate receptors antagonize sensory synaptic response to noxious stimuli of thalamic neurons (Eaton, S. A. et al., *Eur. J. Neuroscience*, 5, 186, 1993).


The use of compounds active at the mGluRs for treatment of neurological diseases such as senile dementia have also been indicated by the findings of Zheng and Gallagher (*Neuron* 9, 163, 1992) and Bashir et al. (*Nature* 363, 347, 1993) who demonstrated that activation of mGluRs is necessary for the induction of long-term potentiation (LTP) in nerve cells (septal nucleus, hippocampus) and the finding that long-term depression is induced after activation of metabotropic glutamate receptors in cerebellar granule cells (Linden et al. *Neuron* 7, 81, 1991).

Thus compounds that demonstrate either activating or inhibiting activity at mGluRs have therapeutic potential for the treatment of neurological disorders. These compounds have application as new drugs to treat both acute and chronic neurological disorders, such as stroke and head injuries; epilepsy; movement disorders associated with Parkinson's disease and Huntington's chorea; pain; anxiety; AIDS dementia; and Alzheimer's disease. Since the mGluRs can influence levels of alertness, attention and cognition; protect nerve cells from excitotoxic damage resulting from ischemia, hypoglycemia and anoxia; modulate the level of neuronal excitation; influence central mechanisms involved in controlling movement; reduce sensitivity to pain; and reduce levels of anxiety, these compounds can also be used to influence these situations and also find use in learning and memory deficiencies such as senile dementia. mGluRs may also be involved in addictive behavior, alcoholism, drug addiction, sensitization and drug withdrawal (*Science*, 280:2045, 1998), so compounds acting at mGluRs might also be used to treat these disorders.

The current pharmaceutical options for treating neurological disorders tend to be very general and non-specific in their actions in that, although they may reduce the clinical symptoms associated with a specific neurological disorder, they may also negatively impact normal function of the central nervous system of patients. Thus new cellular targets and drugs that are more specific in their actions require to be identified and developed and thus a need remains for chemical compounds that demonstrate specific binding characteristics towards mGluRs.

SUMMARY OF THE INVENTION

It is an object of this invention to provide novel compounds that demonstrate activity at the various metabotropic glutamate receptors (mGluRs). In particular, a compound of Formula I and stereoisomers thereof:



wherein:

R1 can be an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol, -CH₂-carboxyl, -CH₂-phosphono, -CH₂-phosphino, -CH₂-sulfono, -CH₂-sulfinio, -CH₂-borono, -CH₂-tetrazol, -CH₂-isoxazol and higher homologues thereof;

R2 can be a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3° amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea;

R3 can be H, aliphatic, aromatic or heterocyclic;

R4 can be an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol;

and pharmaceutically acceptable salts thereof.

DETAILED DESCRIPTION OF THE INVENTION

The terms and abbreviations used in the instant examples have their normal meanings unless otherwise designated. For example "°C" refers to degrees Celsius; "N" refers to normal or normality; "mmol" refers to millimole or millimoles; "g" refers to gram or grams; "mL" means milliliter or milliliters; "M" refers to molar or molarity; "MS" refers to mass spectrometry; "IR"

refers to infrared spectroscopy; and "NMR" refers to nuclear magnetic resonance spectroscopy.


As would be understood by the skilled artisan throughout the synthesis of the compounds of Formula I, it may be necessary to employ an amino-protecting group or a carboxy-protecting group in order to reversibly preserve a reactively susceptible amino or carboxy functionality while reacting other functional groups on the compound

Examples of such amino-protecting groups include formyl, trityl, phthalimido, trichloroacetyl, chloroacetyl, bromoacetyl, iodoacetyl, and urethane-type blocking groups such as benzyloxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, *t*-butoxycarbonyl, 2-(4-xenyl)-isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)-prop-2-yloxycarbonyl, cyclopentanyloxy-carbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)-ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)-ethoxycarbonyl, fluorenylmethoxycarbonyl ("Fmoc"), 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl, 1-piperidyloxycarbonyl and the like; benzoylmethylsulfonyl group, 2-nitrophenylsulfonyl, diphenylphosphine oxide and like amino-protecting groups. The species of amino-protecting group employed is not critical so long as the derivatized amino group is stable to the condition of subsequent reaction(s) on other positions of the intermediate molecule and can be selectively removed at the appropriate point without disrupting the remainder of the molecule including any other amino-protecting group(s). Preferred amino-protecting groups are *t*-butoxycarbonyl (*t*-Boc), allyloxycarbonyl and benzyloxycarbonyl (CbZ). Further examples of these groups are found in E. Haslam in *Protective Groups in Organic Synthesis*; McOmie, J. G. W., Ed. 1973, at Chapter 2; and Greene, T.W. and Wuts, P. G. M., *Protective Groups in Organic Synthesis*, Second edition; Wiley-Interscience: 1991; Chapter 7.

Examples of such carboxyl-protecting groups include methyl, *p*-nitrobenzyl, *p*-methylbenzyl, *p*-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, pentamethylbenzyl, 3,4-methylenedioxybenzyl, benzhydryl, 4,4'-dimethoxybenzhydryl, 2,2',4,4'-tetramethoxybenzhydryl, *t*-butyl, *t*-amyl, trityl,

4-methoxytrityl, 4,4'-dimethoxytrityl, 4,4',4''-trimethoxytrityl, 2-phenylprop-2-yl, trimethylsilyl, *t*-butyldimethylsilyl, phenacyl, 2,2,2-trichloroethyl, β -(di(*n*-butyl)methylsilyl)ethyl, *p*-toluenesulfonyl, 4-nitrobenzylsulfonyl, allyl, cinnamyl, 1-(trimethylsilylmethyl)prop-1-en-3-yl and like moieties. Preferred carboxyl-protecting groups are allyl, benzyl and *t*-butyl. Further examples of these groups are found in E. Haslam, *supra*, at Chapter 5; and T. W. Greene and P. G. M. Wuts, *supra*, at Chapter 5.

The present invention provides a compound of the formula:



wherein:

R1 can be an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol, -CH₂-carboxyl, -CH₂-phosphono, -CH₂-phosphino, -CH₂-sulfono, -CH₂-sulfino, -CH₂-borono, -CH₂-tetrazol, -CH₂-isoxazol and higher analogues thereof;


R2 can be a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3° amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea,

R3 can be H, aliphatic, aromatic or heterocyclic;

R4 can be an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfino, borono, tetrazol, isoxazol;

and pharmaceutically acceptable salts thereof.

In particular compounds wherein the compound of Formula I is selected from the group consisting of:



wherein:

R₁ is COOH

R₂ is COOH

R₃ can be H or methyl or xanthyl or thioxanthyl and

R₄ is NH₂

While all of the compounds of Formula I are believed to demonstrate activity at the metabotropic glutamate receptors (mGluRs), certain groups of Formula I compounds are more preferred for such use.

As noted above, this invention includes the pharmaceutically acceptable salts of the compounds defined by Formula I. A compound of this invention can possess a sufficiently acidic, a sufficiently basic, or both functional groups, and accordingly react with any of a number of organic and inorganic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt.

The term "pharmaceutically acceptable salt" as used herein, refers to salts of the compounds of the above formula which are substantially non-toxic to living organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a pharmaceutically acceptable mineral or organic acid or an organic or inorganic base. Such salts are known as acid addition and base addition salts.

Acids commonly employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydriodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as *p*-toluenesulfonic acid, methanesulfonic acid, oxalic acid, *p*-bromophenylsulfonic acid, carbonic

acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, hydrochloride, dihydrochloride, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, hydroxybenzoate, methoxybenzoate, phthalate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like. Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic acids such as maleic acid and methanesulfonic acid.

Salts of amine groups may also comprise quarternary ammonium salts in which the amino nitrogen carries a suitable organic group such as an alkyl, alkenyl, alkynyl, or aralkyl moiety.

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium hydroxide, calcium carbonate, and the like. The potassium and sodium salt forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is usually not of a critical nature, so long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole. This invention further encompasses the pharmaceutically acceptable solvates of the compounds of Formula I. Many of the Formula I compounds can combine with solvents such as water, methanol, ethanol and acetonitrile to form pharmaceutically acceptable solvates such as the corresponding hydrate, methanolate, ethanolate and acetonitrilate.

The compounds of the present invention have multiple asymmetric (chiral) centers. As a consequence of these chiral centers, the compounds of the present invention occur as racemates, mixtures of enantiomers and as individual enantiomers, as well as diastereomers and mixtures of

diastereomers. All asymmetric forms, individual isomers and combinations thereof, are within the scope of the present invention.

The prefixes "*R*" and "*S*" are used herein as commonly used in organic chemistry to denote the absolute configuration of a chiral center, according to the Cahn-Ingold-Prelog system. The stereochemical descriptor *R* (*rectus*) refers to that configuration of a chiral center with a clockwise relationship of groups tracing the path from highest to second-lowest priorities when viewed from the side opposite to that of the lowest priority group. The stereochemical descriptor *S* (*sinister*) refers to that configuration of a chiral center with a counterclockwise relationship of groups tracing the path from highest to second-lowest priority when viewed from the side opposite to the lowest priority group. The priority of groups is decided using sequence rules as described by Cahn et al., *Angew. Chem.*, 78, 413-447, 1966 and Prelog, V. and Helmchen, G.; *Angew. Chem. Int. Ed. Eng.*, 21, 567-583, 1982)


In addition to the *R,S* system used to designate the absolute configuration of a chiral center, the older D-L system is also used in this document to denote relative configuration, especially with reference to amino acids and amino acid derivatives. In this system a Fischer projection of the compound is oriented so that carbon-1 of the parent chain is at the top. The prefix "D" is used to represent the relative configuration of the isomer in which the functional (determining) group is on the right side of the carbon atom at the chiral center and "L", that of the isomer in which it is on the left.

As would be expected, the stereochemistry of the Formula I compounds is critical to their potency as agonists or antagonists. The relative stereochemistry is established early during synthesis, which avoids subsequent stereoisomer separation problems later in the process. Further manipulation of the molecules then employs stereospecific procedures so as to maintain the preferred chirality. The preferred methods of this invention are the methods employing those preferred compounds.

Non-toxic metabolically-labile esters and amides of compounds of Formula I are ester or amide derivatives of compounds of Formula I that are hydrolyzed in vivo to afford said compounds of Formula I and a pharmaceutically acceptable alcohol or amine. Examples of metabolically-labile esters include esters formed with (1-6C) alkanols in which the alkanol moiety may be optionally substituted by a (1-8C) alkoxy group, for example methanol, ethanol, propanol and methoxyethanol. Examples of metabolically-labile amides include amides formed with amines such as methylamine.


According to another aspect, the present invention provides a process for the preparation of a compound of Formula I, or a pharmaceutically acceptable metabolically-labile ester or amide thereof, or a pharmaceutically acceptable salt thereof, which comprises:

(a) hydrolyzing a compound of formula:




in which **R1** is defined as above, **R5** represents a hydrogen atom or an acyl group and **R4** has the meaning defined above. Preferred values for **R5** are hydrogen and (2-6C) alkanoyl groups, such as acetyl.

(b) hydrolyzing a compound of formula:



in which **R6** and **R7** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof; or:

(c) deprotecting a compound of formula:



in which **R8** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R9** represents a hydrogen atom or a nitrogen protecting group;

whereafter, if necessary and/or desired:

- (i) resolving the compound of Formula I;
- (ii) converting the compound of Formula I into a non-toxic metabolically-labile ester or amide thereof;
- and/or;
- (iii) converting the compound of Formula I or a non-toxic metabolically-labile ester or amide thereof into a pharmaceutically acceptable salt thereof.

The protection of carboxylic acid and amine groups is generally described in McOmie, "Protecting Groups in Organic Chemistry, Plenum Press, NY, 1973, and Greene and Wuts, Protecting Groups in Organic Synthesis, 2nd. Ed., John Wiley & Sons, NY, 1991. Examples of carboxyl protecting groups include alkyl groups such as methyl, ethyl, *t*-butyl and *t*-amyl; aralkyl groups such as benzyl, 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, benzhydryl and trityl; silyl groups such as trimethylsilyl and *t*-butyldimethylsilyl; and allyl groups such as allyl and 1-(trimethylsilylmethyl)prop-1-en-3-yl..

Examples of amine-protecting groups include acyl groups, such as groups of formula **R9** CO in which **R9** represents (1-6C) alkyl, (3-10C) cycloalkyl, phenyl(1-6C) alkyl, phenyl(1-6C) alkoxy, or a (3-10C) cycloalkoxy, wherein a phenyl group may optionally be substituted by one or two


substituents independently selected from amino, hydroxy, nitro, halogeno, (1-6C) alkyl, (1-6C) alkoxy, carboxyl, (1-6C) alkoxycarbonyl, carbamoyl, (1-6C) alkanoylamino, (1-6C) alkylsulphonylamino, phenylsulphonylamino, toluenesulphonylamino, and (1-6C) fluoroalkyl.

The compounds of Formula II are conveniently hydrolyzed in the presence of an acid, such as hydrochloric acid or sulfuric acid, or a base, such as an alkali metal hydroxide, for example sodium hydroxide. The hydrolysis is conveniently performed in an aqueous solvent such as water and at a temperature in the range of 50 to 200 °C.

The compounds of Formula III are conveniently hydrolyzed in the presence of a base, for example an alkali metal hydroxide such as lithium, sodium or potassium hydroxide, or an alkaline earth metal hydroxide such as barium hydroxide. Suitable reaction media include water. The temperature is conveniently in the range of from 50 to 150 °C.

The compounds of Formula IV may be deprotected by a conventional method. Thus, an alkyl carboxyl protecting group may be removed by hydrolysis. The hydrolysis may conveniently be performed by heating the compound of Formula V in the presence of either a base, for example an alkali metal hydroxide such as lithium, sodium or potassium hydroxide, or an alkaline metal hydroxide, such as barium hydroxide, or an acid such as hydrochloric acid. The hydrolysis is conveniently performed at a temperature in the range from 10 to 300 °C. An aralkyl carboxyl protecting group may conveniently be removed by hydrogenolysis. The hydrogenolysis may conveniently be effected by reacting the compound of Formula V with hydrogen in the presence of a Group VIII metal catalyst, for example a palladium catalyst such as palladium on charcoal. Suitable solvents for the reaction include alcohols such as ethanol. The reaction is conveniently performed at a temperature in the range from 0 to 100 °C. An acyl, amine protecting group is also conveniently removed by hydrolysis, for example as described for the removal of an alkyl carboxyl protecting group.

The compounds of Formula II may be prepared by reacting a compound of formula:




with an alkali metal cyanide, such as lithium, sodium or potassium cyanide, and an ammonium halide, such as ammonium chloride, conveniently in the presence of ultrasound. Thus, the ammonium halide is mixed with chromatography grade alumina in the presence of a suitable diluent such as acetonitrile. The mixture is then irradiated with ultrasound, whereafter the compound of Formula V is added, and the mixture is again irradiated. The alkali metal cyanide is then added, followed by further irradiation with ultrasound.

Individual isomers of compounds of Formula II may be made by reacting a compound of the Formula V with the stereoisomers of the chiral agent (*S*)- and (*R*)-phenylglycinol and a reactive cyanide such as trimethylsilyl cyanide.


The compounds of Formula III may be prepared by reacting a compound of Formula V with an alkali metal cyanide, such as lithium, sodium or potassium cyanide, and ammonium carbonate or ammonium carbamate. Convenient solvents include water, dilute ammonium hydroxide, alcohols such as methanol, aqueous methanol and aqueous ethanol. Conveniently the reaction is performed at a temperature in the range of from 10 to 150 °C. If desired, the compounds of Formula III may then be alkylated, for example using an appropriate compound of formula **R6 Cl** and/or **R7 Cl**.

The compounds of Formula V can be prepared by reacting a compound of formula.




with a chlorinating agent such as thionyl chloride or phosphorous(V) chloride, followed by reaction with **R4X** wherein R4 has the meaning defined above and X is halogen or OH.

The compounds of Formula V can also be prepared by oxidizing a compound of formula



under Swern conditions.

The compounds of Formula VI can be prepared from compounds of formula:



by reduction.

If **R1** is CO₂Me, this compound can be bought commercially. If **R1** is another substituent, the compound of Formula VIII can be made using standard procedures.

Many of the intermediates described herein, for example the compounds of Formula II, III and IV are believed to be novel, and are provided as further aspects of the invention.

The Formula I compounds of the present invention are agonists or antagonists at certain metabotropic excitatory amino acid receptors (mGluRs). Therefore, another aspect of the present invention is a method of affecting mGluRs in mammals, which comprises administering to a mammal requiring modulated excitatory amino acid neurotransmission a pharmacologically-effective amount of a compound of Formula I. The term "pharmacologically-effective amount" is used to represent an amount of the compound of the invention that is capable of affecting the mGluRs. By affecting, a compound of the invention is acting as an agonist or antagonist. When a compound of the

invention acts as an agonist, the interaction of the compound with the excitatory amino acid receptor mimics the response of the interaction of this receptor with its natural ligand (i.e. L-Glutamic acid). When a compound of the invention acts as an antagonist, the interaction of the compound with the excitatory amino acid receptor blocks the response of the interaction of this receptor with its natural ligand (i.e. L-Glutamic acid).

The particular dose of compound administered according to this invention will, of course, be determined by the particular circumstances surrounding the case, including the compound administered, the route of administration, the particular condition being treated, and similar considerations. The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, or intranasal routes. Alternatively, the compound may be administered by continuous infusion. A typical daily dose will contain from about 0.001 mg/kg to about 100 mg/kg of the active compound of this invention. Preferably, daily doses will be about 0.05 mg/kg to about 50 mg/kg, more preferably from about 0.1 mg/kg to about 20 mg/kg.

A variety of physiological functions have been shown to be subject to influence by excessive or inappropriate stimulation of excitatory amino acid transmission. The Formula I compounds of the present invention are believed (through their interactions at the mGluRs) to have the ability to treat a variety of neurological disorders in mammals associated with this condition, including acute neurological disorders such as cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia (e.g. stroke and cardiac arrest), spinal cord trauma, head trauma, perinatal hypoxia, and hypoglycemic neuronal damage. The Formula I compounds are believed to have the ability to treat a variety of chronic neurological disorders, such as Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, AIDS-induced dementia, ocular damage and retinopathy, cognitive disorders, and idiopathic and drug-induced Parkinson's disease. The present invention also provides methods for treating these disorders which comprises administering to a patient in need thereof an effective amount of a compound of Formula I.

The Formula I compounds of the present invention (through their interactions at the mGluRs) are also believed to have the ability to treat a variety of other neurological disorders in mammals that are associated with glutamate dysfunction, including muscular spasms, convulsions, migraine headaches, urinary incontinence, psychosis, drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol), smoking cessation, anxiety and related disorders (e.g. panic attack), emesis, brain edema, chronic pain, sleep disorders, Tourette's syndrome, attention

deficit disorder, and tardive dyskinesia. Therefore, the present invention also provides methods for treating these disorders which comprise administering to a patient in need thereof an effective amount of the compound of Formula I.

The Formula I compounds of the present invention (through their interactions at the mGluRs) are also believed to have the ability to treat a variety of psychiatric disorders, such as schizophrenia, anxiety and related disorders (e.g. panic attack), depression, bipolar disorders, psychosis, and obsessive compulsive disorders. The present invention also provides methods for treating these disorders which comprises administering to a patient in need thereof an effective amount of a compound of Formula I.

The pharmacological properties of the compounds of the invention can be illustrated by determining their effects in various functional in vitro assays. The compounds of the invention were studied in an in vitro assay that measured the inhibition of PI hydrolysis or the formation of cyclic AMP in Chinese hamster ovary cell lines expressing mGluR_{1a}, mGluR₂ and mGluR_{4a} cloned metabotropic glutamate receptors.

Principle

So far eight different clones of the G-protein-coupled mGluRs have been identified (Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426). These receptors function to modulate the presynaptic release of L-Glutamate, and the postsynaptic sensitivity of the neuronal cell to L-Glutamate excitation. Based on pharmacology, sequence homology and the signal transduction pathway that they activate, the mGluRs have been subclassified into three groups. The mGluR₁ and mGluR₅ receptors form group I. They are coupled to hydrolysis of phosphatidylinositol (PI) and are selectively activated by (RS)-3,5-dihydroxyphenylglycine (Brabet et al., *Neuropharmacology*, 34, 895-903, 1995). Group II comprises mGluR₂ and mGluR₃ receptors. They are negatively coupled to adenylate cyclase and are selectively activated by (2S,1'R,2'R,3'R)-2-(2,3-dicarboxycyclopropyl)glycine (DCG-IV; Hayashi et al., *Nature*, 366, 687-690, 1993). Finally, the mGluR₄, mGluR₆, mGluR₇ and mGluR₈ receptors belong to group III. They are also negatively coupled to adenylate cyclase and are selectively activated by (S)-2-amino-4-phosphonylbutyric acid (L-AP4; Knopfel et al., 1995, *J. Med. Chem.*, 38, 1417-1426).

Cell Culture

The Chinese hamster ovary cell lines expressing mGluR_{1a}, mGluR₂ and mGluR_{4a} receptors have been described previously (Aramori and Nakanishi, *Neuron* 8, 757-765; 1992; Tanabe et al., *Neuron* 8, 169-179, 1992; Tanabe et al., *J. Neurosci.* 13, 1372-1378). They were maintained at 37°C in a humidified 5% CO₂ incubator in Dulbecco's Modified Eagle Medium (DMEM) containing a reduced concentration of (S)-glutamine (2mM) and were supplemented with 1% proline, penicillin (100 U/ml), streptomycin (100 mg/ml) and 10% dialyzed fetal calf serum (all GIBCO, Paisley). Two days before assay 1.8×10^6 cells were divided into the wells of 24 well plates.

Second Messenger Assays

PI hydrolysis was measured as described previously (Hayashi et al., *Br. J. Pharmacol.* 107, 539-543, 1992; Hayashi et al., *J. Neurosci.* 14, 3370-3377, 1994). Briefly, the cells were labeled with [³H]inositol (2 µCi/ml) 24 h prior to the assay. For agonist assays, the cells were incubated with ligand dissolved in phosphate-buffered saline (PBS)-LiCl for 20 min, and agonist activity was determined by measurement of the level of ³H-labeled mono-, bis- and tris-inositol phosphates by ion-exchange chromatography. For antagonist assays, the cells were preincubated with the ligand dissolved in PBS-LiCl for 20 min prior to incubation with ligand and 10 µM (L)-Glutamic acid for 20 min. The antagonist activity was then determined as the inhibitory effect of the (L)-Glutamic acid-mediated response. The assay of cyclic AMP formation was performed as described previously (Hayashi et al., *Br. J. Pharmacol.* 107, 539-543, 1992; Hayashi et al., *J. Neurosci.* 14, 3370-3377, 1994). Briefly, the cells were incubated for 10 min in PBS containing the ligand and 10 µM forskolin and 1mM 3-Isobutyl-1-methyloxanthine (IBMX; both Sigma, St. Louis, MO, USA). The agonist activity was then determined as the inhibitory effect of the forskolin-induced cyclic AMP formation. For antagonist assay, the cells were preincubated with ligand dissolved in PBS containing 1 mM IBMX for 20 min prior to a 10 min incubation in PBS containing the ligand, 20 µM (mGluR₂) or 50 µM (mGluR_{4a}), (L)-Glutamic acid, 10 µM Forskolin and 1 mM IBMX.

Results

Some of the compounds of the invention were tested for antagonist activity against Chinese hamster

ovary cell lines expressing mGluR_{1α}, mGluR₂ and mGluR_{4α} cloned mGluRs at a concentration of 1 mM. When tested as antagonists of the increase in PI hydrolysis evoked by 10 μM (L)-Glutamic acid, some compounds of the invention effectively blocked this increase in PI hydrolysis by an action at the mGluR_{1α} receptor. The data for one of the compounds of the invention is shown in Figure 1 below

According to another aspect, the present invention provides a method of modulating one or more metabotropic glutamate receptor functions in a warm-blooded mammal which comprises administering an effective amount of a compound of Formula I, or a non-toxic metabolically-labile ester or amide thereof, or a pharmaceutically acceptable salt thereof.

The compounds of the present invention are preferably formulated prior to administration. Therefore, another aspect of the present invention is a pharmaceutical formulation comprising a compound of Formula I and a pharmaceutically-acceptable carrier, diluent, or excipient. The present pharmaceutical formulations are prepared by known procedures using well-known and readily available ingredients. In making the compositions of the present invention, the active ingredient will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier, and may be in the form of a capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be a solid, semi-solid, or liquid material that acts as a vehicle, excipient, or medium for the active ingredient.

The compounds of Formula I are usually administered in the form of pharmaceutical compositions. These compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. These compounds are effective as both injectable and oral compositions. Such compositions are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound.

The present invention also provides pharmaceutical compositions containing compounds as disclosed in the claims in combination with one or more pharmaceutically acceptable, inert or physiologically active, diluent or adjuvant. The compounds of the invention can be freeze-dried and, if desired, combined with other pharmaceutically acceptable excipients to prepare formulations for administration. These compositions may be presented in any form appropriate for the administration route envisaged. The parenteral and the intravenous route are the preferential routes for administration.

Compounds of the general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs.

Compositions intended for oral use may be prepared according to any known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium

carboxymethylcellulose, methyl cellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example hepta-decaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or *n*-propyl-*p*-hydroxy benzoate, one or more colouring agents, one or more flavouring agents or one or more sweetening agents, such as sucrose or saccharin

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example peanut oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin.

The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavouring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavouring and colouring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oil phase may be a vegetable oil, for example olive oil or peanut oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavouring and colouring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or a suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compound(s) of the general Formula I may be administered, together or separately, in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compound(s) of general Formula I may be administered, together or separately, parenterally in sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anaesthetics, preservatives and buffering agents can be dissolved in the vehicle.

The dosage to be administered is not subject to defined limits, but it will usually be an effective amount. It will usually be the equivalent, on a molar basis of the pharmacologically active free form produced from a dosage formulation upon the metabolic release of the active free drug to achieve its desired pharmacological and physiological effects. The compositions are preferably formulated in a unit dosage form, each dosage containing from about 0.05 to about 100 mg, more usually about 1.0 to about 30 mg, of the active ingredient. The term "unit dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

The active compound is effective over a wide dosage range. For examples, dosages per day normally fall within the range of about 0.01 to about 30 mg/kg of body weight. A typical daily dose will contain from about 0.01 mg/kg to about 100 mg/kg of the active compound of this invention. Preferably, daily doses will be about 0.05 mg/kg to about 50 mg/kg, more preferably from about 0.1 mg/kg to about 25 mg/kg. In the treatment of adult humans, the range of about 0.1 to about 15 mg/kg/day, in single or divided dose, is especially preferred. However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. In some instances dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several smaller doses for administration throughout the day.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 5 mg to about 500 mg, more preferably about 25 mg to about 300 mg of the active ingredient. The term "unit dosage form" refers to a physically discrete unit suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier, diluent, or excipient. The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way.

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

	Quantity (mg/capsule)
Active Ingredient	250
Starch, dried	200
Magnesium stearate	10
Total	460

The above ingredients are mixed and filled into hard gelatin capsules in 460 mg quantities

Formulation 2

A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
Active Ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	5
Total	665

The components are blended and compressed to form tablets each weighing 665 mg.

Formulation 3

An aerosol solution is prepared containing the following components:

	Weight %
Active Ingredient	0.25
Ethanol	29.75
Propellant 22 (Chlorodifluoromethane)	70.00
Total	100

The active compound is mixed with ethanol and the mixture added to a portion of the Propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

Formulation 4

Tablets each containing 60 mg of active ingredient are made as follows:

	Quantity (mg/tablet)
Active Ingredient	60
Starch	45
Microcrystalline cellulose	35
Polyvinylpyrrolidone	4
Sodium carboxymethyl starch	4.5
Magnesium stearate	0.5
Talc	1.0
Total	150

The active ingredient, starch, and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders that are then passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and passed

through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

Formulation 5

Capsules each containing 80 mg medicament are made as follows.

	Quantity (mg/capsule)
Active Ingredient	80
Starch	59
Microcrystalline cellulose	59
Magnesium stearate	2
Total	200

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 sieve, and filled into hard gelatin capsules in 200 mg quantities.

Formulation 6

Suppositories each containing 225 mg of active ingredient may be made as follows:

	Quantity (mg/suppository)
Active Ingredient	225
Saturated fatty acid glycerides	2000
Total	2225

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

Formulation 7

Suspensions each containing 50 mg of medicament per 5 mL dose are made as follows:

Active Ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mL
Benzoic acid solution	0.10 mL
Flavour	q.v.
Color	q.v.
Purified water to total	5 mL

The medicament is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with some of the water and added, with stirring. Sufficient water is then added to produce the required volume.

Formulation 8

An intravenous formulation may be prepared as follows:

	Quantity
Active Ingredient	100 mg
Mannitol	100 mg
5 N Sodium hydroxide	200 mL
Purified water to total	5 mL

Formulation 9

A topical formulation may be prepared as follows:

	Quantity
Active Ingredient	1-10 g
Emulsifying Wax	30 g
Liquid Paraffin	20 g
White soft paraffin to	100 g

The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

Formulation 10

Sublingual or buccal tablets, each containing 10 mg of active ingredient, may be prepared as follows:

	Quantity (mg/tablet)
Active Ingredient	10.0
Glycerol	210.5
Water	143.0
Sodium Citrate	4.5
Polyvinyl Alcohol	26.5
Polyvinylpyrrolidone	15.5
Total	410.0

The glycerol, water, sodium citrate, polyvinyl alcohol, and polyvinylpyrrolidone are admixed together by continuous stirring and maintaining the temperature at about 90 °C. When the polymers have gone into solution, the solution is cooled to about 50°-55 °C and the medicament is slowly admixed. The homogenous mixture is poured into forms made of an inert material to produce a drug-containing diffusion matrix having a thickness of about 2-4 mm. This diffusion matrix is then cut to form individual tablets having the appropriate size.

Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in controlled amounts

The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art (see, for example, U.S. Pat. No. 5,023,252, issued Jun. 11, 1991) herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.


Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of biological factors to specific anatomical regions of the body, is described in U.S. Pat. No. 5,011,472, issued Apr. 30, 1991, which is herein incorporated by reference.

Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs or prodrugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions that can transiently open the blood-brain barrier.

EXAMPLES

The following Examples illustrate the invention. The following abbreviations are used in the Examples: EtOAc, ethyl acetate; THF, tetrahydrofuran; EtOH, ethanol; TLC, thin layer chromatography; GC, gas chromatography; HPLC, high pressure liquid chromatography; m-CPBA, m-chloroperbenzoic acid; Et₂O, diethyl ether; DMSO, dimethyl sulfoxide; DBU, 1,8-diazabicyclo-[5.4.0]undec-7-ene, MTBE, methyl *t*-butyl ether; FDMS, field desorption mass spectrometry and r.t., room temperature.

Example 1: Synthesis of Cubanylglycinates IGT 1.0 series

*Preparation 1: 4-methoxycarbonylcubane carboxylic acid*

A solution of cubane dimethyl ester (6.0g, 27.24 mmol) in 182 mL of dry THF is stirred under N_2 at room temperature. A solution of methanolic NaOH (26.7 mmol, 10.7 mL 2.5 M) is added dropwise from a pressure equalized addition funnel and the resulting solution stirred at room temperature for 16 h. The mixture is evaporated under reduced pressure at r.t., the residue is taken up in 66 mL of water and extracted with 3 x 25 mL of chloroform. The aqueous layer is acidified to pH 3 with concentrated HCl and extracted with 3 x 30 mL of chloroform. The combined organic layers were dried over magnesium sulphate, filtered and evaporated to give (2) 182-183 °C: 1H NMR ($CDCl_3$) δ 3.72 (s, 3H), 4.27 (m, 6H).
Yield 5.1 g (91%).

Preparation 2: 4-methoxycarbonyl-1-(hydroxymethyl) cubane

The mono acid (2) (0.48 g) is dissolved in dry THF (5 mL) and cooled to -70 °C. A solution of BH_3 in THF is added slowly with stirring. The reaction mixture is stirred at -78 °C for 4 hrs and

allowed to come to room temperature. Water (3 mL) is added and stirred for 30 min, potassium carbonate (0.85 g) is added and the solution extracted with Et₂O. The organic phase is dried over magnesium sulfate and evaporated to give the alcohol (3) 0.46 g (100%) m.p. 83-85 °C. ¹H NMR(200 MHz, solvent) δ: 1.58 (s, 1H), 3.62 (s, 3H), 3.72 (s, 2H), 3.81 (m, 3H), 4.1 (m, 3H).

Preparation 3: 4-methoxycarbonyl-1-(formyl) cubane

DMSO (0.7 mL, 9.68 mmol) is added to oxalyl chloride (0.42 mL, 4.84 mmol) in 12 mL of CH₂Cl₂ at -78 °C. The alcohol (3) (0.46 g, 2.42 mmol) in 3 mL CH₂Cl₂ is added and stirred at -78 °C for 1.5 h. Triethylamine (2.0 mL, 14.4 mmol) is added and the mixture is allowed to come to 0 °C. Saturated ammonium chloride solution is added and the phases separated, the aqueous layer is extracted with CH₂Cl₂ and the combined organic layers are dried (MgSO₄), then evaporated to give crude product which is purified by flash chromatography (1:1 hexanes:diethyl ether) to give 0.35 g (76%) of pure product (4). ¹H NMR (200 MHz, solvent) δ: 3.7 (s, 3H), 4.2 (m, 3H), 4.32 (m, 3H), 9.72 (s, 1H).


Preparation 4: 4-methoxycarbonyl-1-[2'-hydroxy-1'-phenylethyl] methylnitrilecubane

(*R*)-phenylglycinol (257 mg, 1.87 mmol) is added to a solution of the aldehyde (4) (0.35 g, 1.84 mmol) in 14 mL of methanol. The solution is cooled to 0 °C and TMSCN (0.49 mL, 3.68 mmol) is added and the mixture stirred at 0 °C overnight. Evaporation of the solvent leaves a residue which is purified by chromatography (diethyl ether:hexanes, 3:1) to give 0.48 g (77%) of pure product (5). ¹H NMR (CDCl₃) δ: 2.23 (s, 1H), 2.6 (br, 1H), 3.5-3.75 (m, 2H), 3.7 (s, 3H), 3.9 (m, 3H), 4.11 (dd, 1H), 4.2 (m, 3H), 7.3 (s, 5H).

Preparation 5: 4-carbonyl-1-cubanylglycine

Lead acetate (0.69 g, 1.57 mmol) is added to a stirred solution of nitrile (5) (0.48 g, 1.42 mmol) in dry methanol/dichloromethane 1:1 (12 mL). After 10 min 10 mL of water is added and the suspension filtered through celite. The organic layer is dried and evaporated to give the crude imine. The crude imine is refluxed with 6N HCl (30 mL) for 6 hr. The solution is evaporated to dryness and placed on anion exchange resin, eluting with 1N acetic acid to yield the product (6). mp. 241 °C (dec.) ¹H NMR (D₂O) δ 3.96 (s, 1H), 4.01 (m, 3H), 4.14 (m, 3H).

Example 2

*Preparation 1: 4-methoxycarbonylcubane carboxylic acid*

A solution of cubane dimethyl ester (6.0g, 27.24 mmol) in 182 mL of dry THF is stirred under N_2 at r.t. a solution of methanolic NaOH (26.7 mmol, 10.7 mL 2.5 M) is added dropwise from a pressure equalized addition funnel and the resulting solution stirred at r.t. for 16 h. The mixture is evaporated under reduced pressure at r.t., the residue is taken up in 66 mL of water and extracted with 3 x 25 mL of chloroform. The aqueous layer is acidified to pH 3 with concentrated HCl and extracted with 3 x 30 mL of chloroform. The combined organic layers were dried over magnesium sulphate, filtered and evaporated to give (2) 182-183 °C: ^1H NMR (CDCl_3) δ 3.72 (s, 3H), 4.27 (m, 6H).

Yield 5.1 g (91%).

Preparation 2: 4-methoxycarbonylcubane-1-carbonyl chloride

The monomethyl ester (2) (1.37 g, 6.65 mmol) is dissolved in 15 mL of thionyl chloride and gently refluxed overnight. The thionyl chloride is evaporated off and the resultant residue containing (3) was used immediately without further purification.

Preparation 3: 4-methoxycarbonylcubane-1-methyl ketone

A suspension of copper iodide (1.49 g, 7.83 mmol) in 30 mL of dry THF is stirred at 0°C. Methyl lithium (15.75 mmol, 11.2 mL of 1.4 M) was added and stirred at 0°C for 30 min, then cooled to -78°C. A solution of 1.6 g, 7.12 mmol of (3) in 10 mL dry THF is added and the resultant mixture stirred for 1 h. at -78°C. The mixture was quenched with saturated ammonium chloride solution (15 mL) and extracted with 3 x 30 mL of diethyl ether. The combined organic layers were dried over magnesium sulphate, filtered and evaporated to give crude (4). The product was purified by silica chromatography (hexanes:ethyl acetate, 2:1) to give 1.0 g of product (yield 69%). m.p. 87-89°C. ^1H NMR (CDCl_3) δ 2.17 (s, 3H), 3.7 (s, 3H), 4.21 (m, 6H).


Preparation 4: 4-methoxycarbonylcubane-1-methyl-1-(5,5'-hydantoin)

A solution of the methyl ketone (4) (1.0 g, 4.9 mmol) in 40 mL of ethanol and 5.8 mL of 1 N NaOH, is stirred at 70°C for 4 h. The resulting solution is evaporated to dryness under reduced pressure and redissolved in 1:1 ethanol: water (20 mL). To this solution is added potassium cyanide (0.35 g, 5.4 mmol) and ammonium carbonate (0.96 g, 9.8 mmol) and the mixture heated in a sealed tube at 85°C for 24 h. The reaction is cooled, acidified with 6 N HCl and reduced in volume until a precipitate forms. The precipitate is filtered and the filtrate evaporated to dryness and extracted with ethyl acetate. The solvent is evaporated and the product combined with the residue from above to give (5) as a white solid. Yield 0.95 g (75%) m.p. 244-248°C. NMR ^1H (DMSO) δ 1.18 (s, 3H) 3.9 (m, 3H), 4.0 (m, 3H), 8.1 (s, 1H), 10.6 (s, 1H).

Preparation 5: 4-carboxycubane-1-methylglycine

The hydantoin (5) (0.95 g, 3.65 mmol) is dissolved in 30 mL of 2 N NaOH and heated to 170°C in a sealed tube for 20 h. The reaction is cooled and filtered to remove precipitate and the filter cake washed with 3 x 10 mL of water. The combined aqueous washings are evaporated to give crude (6) which is applied to Spectrum 1X4 anion exchange resin, eluted with 0.5 N acetic acid. Isolation by evaporation and crystallization gives (6) as colorless crystals. m.p. >250°C (decomp.). NMR. ^1H (D_2O) δ 1.38 (s, 3H), 3.95 (s, 6H).

Example 3

*Preparation 1: 4-methoxycarbonylcubane-1-acyl ethylcarboxylate.*

n-butyl lithium (34.83 mmol, 23.5 mL of 1.5 M) is added dropwise to a stirred solution of ethyl hydrogen malonate (2.32 g, 17.41 mmol) in 80 mL of dry THF under N₂ at -78°C. The mixture was warmed to -30°C over 0.5 h and then re-cooled to -78°C. The acid chloride of cubane monomethyl ester from example (2) above (2.35 g, 10.46 mmol) in 7 mL of THF is added dropwise to the stirred solution. The reaction is warmed slowly to r.t and stirred for a further 1 h.

The solution is poured into 50 mL of 1 N HCl and extracted with 3 x 50 mL of diethyl ether.

The combined organic extracts are further extracted with 20 mL of saturated sodium hydrogen carbonate and brine, dried over magnesium sulphate, filtered and evaporated to give crude (2).

The product is purified by column chromatography on silica with hexanes: ethyl acetate 2:1 to yield 2.5 g (86%) of (2). ¹H NMR (CDCl₃) δ 1.2 (t, 3H) 3.4 (s, 2H), 3.65 (s, 3H), 4.2 (m, 8H).

Preparation 2: 4-methoxycarbonylcubane-1-(thioxanthyl)-acetyl ethylcarboxylate.

cubane- β -ketoester (2) (1.15g, 4.16 mmol) and thioxanthene-9-ol (0.88g, 4.1 mmol) are dissolved in 18 mL of a 1:1 mixture of ethanol:acetic acid and stirred at r.t. for 3 days. The resulting crystalline solid was filtered off to yield 1.52 g (77%) of pure (3) m.p. 147-149°C. ^1H NMR (CDCl_3) 1.00 (t, 3H), 3.24 (s, 3H), 3.75 (m, 3H), 3.9 (q, 2H), 4.0 (m, 3H), 4.6 (d, 1H), 5.0 (d, 1H), 7.3 (m, 8H).

Preparation 3: 4-carboxycubane-1-methylthioxanthylketone

The thioxanthylcubane adduct (3) (1.69 g, 3.57 mmol) is dissolved in ethanol 33 mL and 8.7 mL of 1 N NaOH and heated at 70°C for 4 h. The resulting solution is evaporated and redissolved in 25 mL of water, acidified with 6 N HCl and extracted with 3 x 50 mL of diethyl ether. The combined organic layers are dried over magnesium sulphate, filtered and concentrated to give a crude product containing (4). Chromatography on silica using ethyl acetate gives 1.26 g (88%) of (4)

^1H NMR (CDCl_3) δ 2.8 (d, 2H), 3.8 (m, 3H), 4.0 (m, 3H), 4.7 (t, 1H), 7.3 (m, 8H), 9.5 (br, 1H).

Preparation 4: 4-carboxycubane-1-thioxanthyl-1-(5,5'-hydantoin)

The thioxanthyl cubane ketone (4) (1.24 g, 3.22 mmol) is dissolved in 1:1 ethanol:water (20 mL) Potassium cyanide (0.522 g, 8.0 mmol) and ammonium carbonate (1.39 g, 14.4 mmol) are added and the solution heated in a sealed tube at 85°C for 65 h. The reaction is cooled and acidified with 2 N HCl and extracted with 3 x 40 mL of ethyl acetate. The organic layers are combined, dried over magnesium sulphate, filtered and evaporated to give (5) 1.3 g (88%) as a crude product. This material was hydrolyzed in the next step without purification

^1H NMR (CD_3OD) δ 1.7 (m, 1H), 2.7 (m, 1H), 3.8 (m, 3H), 4.0 (m, 3H), 4.3 (m, 1H), 7.4 (m, 8H).

Preparation 5: 4-carboxycubane-1- thioxanthyl lglycine


The hydantoin adduct (5) (300 mg, 0.65 mmol) is taken up in 1 N NaOH (10 mL) and heated at 170 °C for 20 h in a sealed tube. The mixture is cooled and the pH adjusted with 6 N HCl to between 7 and 8. The precipitate formed is filtered and washed with water. The combined filtrate and washings are combined and evaporated to dryness. The resulting residue is purified by column chromatography and finally by reverse phase chromatography to yield (6) as colorless crystals. 70 mg. ^1H NMR ($\text{CD}_3\text{OD} + \text{D}_2\text{O}$) δ 2.3 (m, 2H), 3.9 (s, 6H), 4.4 (m, 1H), 7.4 (m, 8H).

20-07-2000

CA 009900311

We claim:

1. A compound of the formula:



wherein:

R1 can be an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol, $-\text{CH}_2\text{-carboxyl}$, $-\text{CH}_2\text{-phosphono}$, $-\text{CH}_2\text{-phosphino}$, $-\text{CH}_2\text{-sulfono}$, $-\text{CH}_2\text{-sulfinio}$, $-\text{CH}_2\text{-borono}$, $-\text{CH}_2\text{-tetrazol}$, and $-\text{CH}_2\text{-isoxazol}$;

R2 can be a basic group selected from the group consisting of 1° amino, 2° amino, 3° amino, quaternary ammonium salts, aliphatic 1° amino, aliphatic 2° amino, aliphatic 3° amino, aliphatic quaternary ammonium salts, aromatic 1° amino, aromatic 2° amino, aromatic 3° amino, aromatic quaternary ammonium salts, imidazol, guanidino, boronoamino, allyl, urea, thiourea,

R3 can be H, aliphatic, aromatic or heterocyclic;


R4 can be an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol; and pharmaceutically acceptable salts thereof.

2. A compound as claimed in claim 1, wherein **R1** is COOH .
3. A compound as claimed in claim 1, wherein **R2** is NH_2 .

20-07-2000


CA 009900311

4. A compound as claimed in claim 1, wherein R_3 can be -H, or -Me; or xanthyl or thioxanthyl or $-\text{CH}_2\text{-xanthyl}$, or $-\text{CH}_2\text{-thioxanthyl}$ and R_4 is $-\text{COOH}$.
5. A process for the preparation of a compound of Formula I, or a pharmaceutically acceptable metabolically-labile ester or amide thereof, or a pharmaceutically acceptable salt thereof, which comprises:
- (a) hydrolyzing a compound of formula:



wherein: R'_1 is an acidic group selected from the group consisting of carboxyl, phosphono, phosphino, sulfono, sulfinio, borono, tetrazol, isoxazol, $-\text{CH}_2\text{-carboxyl}$, $-\text{CH}_2\text{-phosphono}$, $-\text{CH}_2\text{-phosphino}$, $-\text{CH}_2\text{-sulfono}$, $-\text{CH}_2\text{-sulfinio}$, $-\text{CH}_2\text{-borono}$, $-\text{CH}_2\text{-tetrazol}$, $-\text{CH}_2\text{-isoxazol}$ and higher analogues thereof, or a protected form thereof, R_3 can be H, aliphatic, aromatic or heterocyclic and [in which R_1 is defined as above,] R_5 represents a hydrogen atom or an acyl group [and R_4 has the meaning defined above,], and wherein [P] preferred values for R_5 are hydrogen and (2-6C) alkanoyl groups, such as acetyl; or


- (b) deprotecting and hydrolyzing a compound of formula (II b)



wherein: R'_1 and R_3 are as defined above; or


20-07-2000

(c) [b] hydrolyzing a compound of formula:



wherein: **R6** and **R7** each independently represent a hydrogen atom, a (2-6C) alkanoyl group, a (1-4C) alkyl group, a (3-4C) alkenyl group or a phenyl (1-4C) alkyl group in which the phenyl is unsubstituted or substituted by halogen, (1-4C) alkyl or (1-4C) alkoxy, or a salt thereof, **R'1** and **R3** are as defined above; or

(d) [c] deprotecting a compound of formula:



wherein: [in which] **R8** represents a hydrogen atom or a carboxyl protecting group, or a salt thereof, and **R9** represents a hydrogen atom or a nitrogen protecting group, **R'1** and **R3** are as defined above;

whereafter, if necessary and/or desired:

- (i) resolving the compound of Formula I;
- (ii) converting the compound of Formula I into a non-toxic metabolically-labile ester or amide thereof; and/or;
- (iii) converting the compound of Formula I or a non-toxic metabolically-labile ester or amide thereof into a pharmaceutically acceptable salt thereof.


6. A pharmaceutical formulation, which comprises a compound as claimed in claim 1 and a pharmaceutically acceptable carrier, diluent or excipient.

20-07-2000

CA 009900311


7. A use of the compound according to claim 1 to modulate [A method of modulating] one or more metabotropic glutamate receptor functions in a warm blooded mammal [requiring such treatment], wherein said use [which] comprises administering an effective amount of a compound of formula (I) as claimed in claim 1.

8. A compound of formula:



wherein: R'1, R3 and R5 have the meanings as defined in claim 5 [in which R1, R4 and R5 have the meanings as defined above].

9. A compound of formula:




wherein: R'1, R3, R6 and R7 have meanings as defined in claim 5 [wherein R6 and R7 have meanings as defined above].

20-07-2000

CA 009900311

10. A compound of formula:




wherein: **R'1**, **R3**, **R8** and **R9** have meanings as defined in claim 5 [in which **R8** and **R9** have meanings as defined above].

11. A compound according to claim 1, wherein **R1** is -COOH, **R2** is -NH₂, **R3** is H and **R4** is COOH.
12. A compound according to claim 1, wherein **R1** is -COOH, **R2** is -NH₂, **R3** is CH₃ and **R4** is COOH.
13. A compound according to claim 1, wherein **R1** is -COOH, **R2** is -NH₂, **R3** is -CH₂-thioxanthy and **R4** is COOH.
14. A use of the compound according to claim 1 for the treatment of a neurological disease or disorder selected from the group comprising: cerebral deficits subsequent to cardiac bypass surgery and grafting, cerebral ischemia, stroke, cardiac arrest, spinal cord trauma, head trauma, perinatal hypoxia, and hypoglycemic neuronal damage, Alzheimer's disease, Huntington's Chorea, amyotrophic lateral sclerosis, AIDS-induced dementia, ocular damage, retinopathy, cognitive disorders, idiopathic and drug-induced Parkinson's disease, muscular spasms, convulsions, migraine headaches, urinary incontinence, psychosis, drug tolerance, withdrawal, and cessation (i.e. opiates, benzodiazepines, nicotine, cocaine, or ethanol), smoking cessation, anxiety and related disorders (e.g. panic attack), emesis, brain edema, chronic pain, sleep disorders, Tourette's syndrome, attention deficit disorder, and tardive dyskinesia, wherein said use comprises administering an effective amount of a compound of formula (I).


20-07-2000

CA 009900311

15. A use of the compound according to claim 1 for the treatment of a psychiatric disease or disorder selected from the group comprising: schizophrenia, anxiety and related disorders (e.g. panic attack), depression, bipolar disorders, psychosis, and obsessive compulsive disorders, wherein said use comprises administering an effective amount of a compound of formula (I).
16. The use according to any one of claims 7, 14 or 15 wherein said compound is selected from the group of compounds comprising :



17. A use of the compound:




for the treatment of cerebral ischemia, stroke and cardiac arrest, wherein said use comprises administering an effective amount of the said compound.

20-07-2000

CA 009900311

18. A compound of formula:



wherein: R'1 and R3 have the meaning as defined in claim 5.


19. A compound according to claim 18, wherein: R'1 is -COOMe, R3 is H.

20. A compound according to claim 9, wherein: R'1 is -COOH, R3 is CH₃, R6 = R7 is H.

21. A compound according to claim 9, wherein: R'1 is -COOH, R3 is -CH₂-thioxanthyl, R6 = R7 is H.

Figure 1

The Actions of a Compound of the Invention as an Antagonist of PI Hydrolysis
evoked through the mGluR1 receptor by 10 μ M (L)-Glu



DECLARATION FOR PATENT APPLICATION

Attorney Docket: 24406
Page 2 of 2

We hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. § 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor: ^{KL} ~~Kenneth CURRY~~ HASSAN PAJOUHESHInventor's Signature *H. Pajouhesh*

Date _____

Residence: 606-1010 Pacific Blvd., Vancouver, British Columbia V6Z 2X8 CANADA CAXCountry of Citizenship: CANADAPost Office Address: same as residenceFull name of second inventor: ^{KL} ~~Hassan PAJOUHESH~~ KENNETH CURRYInventor's Signature *Kenneth Curry*

Date _____

Residence: 2438 West 49th Avenue, Vancouver, British Columbia V6M 2V3 CANADA CAXCountry of Citizenship: IRPost Office Address: same as residence

DECLARATION FOR PATENT APPLICATION

Attorney Docket: 24406

Page 1 of 2

As a below-named inventor(s), I/we hereby declare that:

My/Our residence(s), post office address(es) and citizenship(s) is/are as stated below next to my/our name(s).

I/We believe I/we am/are the original inventor, first and sole (if only one name is listed below) or the original, first and joint inventors (if plural names are listed below) of the subject matter which is claimed, and for which a patent is sought on the invention entitled:

CUBANE DERIVATIVES AS METABOTROPIC GLUTAMATE RECEPTOR AGONISTS OR ANTAGONISTS AND PROCESS FOR THEIR PREPARATION

the specification of which: (check one)

☐ is attached hereto.

☒ was filed on 19 April 1999, as Serial No. PCT/CA99/00311,
16 October 2000 09/673,473
 and was amended on _____ (if applicable).

We hereby state that we have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We acknowledge the duty to disclose information which is material to the patentability of this application as defined by 37 CFR § 1.56.

We hereby claim foreign priority benefits under 35 U.S.C. § 119 of any foreign application(s) for patent or inventor's certificate listed below, and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Applications:

			Priority Claimed	
<u>2,235,119</u>	<u>CA</u>	<u>17 / April / 1998</u>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
(Application No.)	(Country)	(Day/Month/Year Filed)	Yes	No
<u> </u>	<u> </u>	<u> / / </u>	<input type="checkbox"/>	<input type="checkbox"/>
(Application No.)	(Country)	(Day/Month/Year Filed)	Yes	No
<u> </u>	<u> </u>	<u> / / </u>	<input type="checkbox"/>	<input type="checkbox"/>
(Application No.)	(Country)	(Day/Month/Year Filed)	Yes	No

We hereby appoint Gary M. Nath, Reg. No. 26,965; Harold L. Novick, Reg. No. 26,011; Todd L. Juneau, Reg. No. 40,669; Lee C. Heiman, Reg. No. 41,827; Jerald L. Meyer, Reg. No. 41,194; Joshua B. Goldberg, Reg. No. 44,126; David R. Murphy, Reg. No. 22,751; Paul A. Sacher, Reg. No. 43,418; Deborah H. Yellin, 45,904; Nahied K. Usman, Reg. No. 47,148; and Roger Hahn, Reg. No. 46,376; as my attorneys to prosecute this application and transact all business in the U.S. Patent and Trademark Office connected therewith.

Direct Telephone Calls to:

Gary M. Nath
(202) 775-8383

Send Correspondence to:
NATH & ASSOCIATES PLLC
Sixth Floor
1030 15th Street, N.W.
Washington, D.C. 20005 U.S.A.

We hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by 35 U.S.C. § 112, first paragraph, I/we acknowledge the duty to disclose material information as defined in 37 CFR § 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

<u>(U.S. Application Serial No.)</u>	<u>(U.S. Filing Date)</u>	<u>(Status--patented, pending, abandoned)</u>
<u>(U.S. Application Serial No.)</u>	<u>(U.S. Filing Date)</u>	<u>(Status--patented, pending, abandoned)</u>